Assessing Global Change Impact on the US using National Lightning Data

Project Status

National Climate Assessment November 20, 2012

(abbreviated version)

William Koshak and Richard Blakeslee Earth Science Office, NASA Marshall Space Flight Center, Huntsville, AL, USA

Project Focus

CLIMATE CHANGE (ΔT)



scientific literature CHANGES IN
US
CLOUD-TO-GROUND
LIGHTNING



NOAA
Storm Data
& NIFC

IMPACTS:

- Human Health
- Agriculture
- Forestry

Assessment Tool

 LSAT software for analyzing NLDN data


Accomplishments


- ☐ Developed a Lightning Software Analysis Tool (LSAT)
 - written in IDL programming language
 - o ingests, calculates, and visualizes national CG lightning data
 - o now serves as a new "sustaining assessment" tool
- □ Applied LSAT to analyze CG lightning over a region slightly larger than CONUS during the period 2003-2011.
- ☐ Used NOAA Storm Data, and National Interagency Fire Center (NIFC) data to obtain lightning-caused death/injury, crop damage, wildfire stats.
- ☐ Compared average values (2003-2006) with average values (2007-2010):
 - ✓ CG lightning frequency dropped by 10.7%
 - ✓ Fatalities dropped by 13.5%
 - ✓ Injuries dropped by 31.2%
 - ✓ Crop damage dropped by 61.25%
 - √ # wildfires dropped by 23.6%
 - ✓ Wildfire burn acreage dropped by 8.3%
 - ✓ Multiplicity dropped by 2.4%
 - ✓ Peak current increased by 9.9%
- □ Number of +CG (and +CG fraction) monotonically trended upward in 2003-2011

Positive CG Fraction Trends Up


(linked to Severe Wx)


Sample LSAT output





POSITIVE CG FRACTION





Accomplishments (cont.)

- ☐ Examined effects of National Lightning Detection Network Upgrades
 - 2002-2003 Upgrade (all sensors replaced w/new IMPACT-ESPs + 8 sensors added)
 - 2004 Propagation Model Upgrade (increases peak current values)
 - 2006 Sensor Addition Upgrade (2 sensors added SE of Florida)
 - 2006 E-Field Waveform Detection Criteria Upgrade (short PTZ waveforms admitted to allow limited IC detection; increases CG count but some are cloud flashes)
 - 2006 15kA Rule Upgrade (no effect since already accounted for)
 - 2008 Location Algorithm Upgrade (extend range to offshore & N. Mexico)
 - 2008 Duplicate/Misplaced Events Upgrade (improvements in removing these)
- □ Only important effect seems to be a network change in 2008 that abruptly increased +CG fraction. Still investigating.

Accomplishments (cont.)

☐ Completed conservative risk-based assessment

Human Health:

Fatalities: 13.98 deaths per 1°C

Injuries: 87.47 injuries per 1°C

Agriculture:

Crop Damage: \$49,348 per 1°C

Forestry:

Wildland Fires (Number): 4091.0 wildfires per 1°C

Wildland Fires (Acres): 936,097.6 acres per 1°C


☐ Completed a 32 page Technical Input Report (TIR) to summarize findings.

☐ Submitted TIR on February 22, 2012. Peer-Reviewed & Revised by May 3.

^{*}Reeve & Toumi (1999): 40±14% change in CG frequency per 1°C; and Price & Rind (1994) found that CG frequency more sensitive to temperature changes than cloud flash frequency.

Plans for Future

- □ Apply LSAT to complete 2012 analyses.
- □ Convert TIR into journal article for publication
 - > Add 2012 results
 - Continue Examining Effects of Network Upgrades on Lightning
 - Examine Natural Effects on Lightning (other than Climate Change):



Plans for Future (cont.)

- ☐ Thru ROSES A.47 Call, develop & track <u>additional indicators</u>:
 - > TRMM/LIS data to examine lightning NOx in Southern CONUS:
 - Total (CG + cloud flash) lightning
 - Flash optical area
 - Flash optical radiance

Increases in these increase NOx & therefore influence
Ozone and OH radical concentrations (hence climate).

- Collaborate with City College of New York & NOAA/CREST to develop & track <u>additional indicators</u> that use CONUS radar data:
 - (Lightning/Rain) Ratio: Important for "dry lightning" (Wildfires)
 - (Lightning x Rain) Product: Characterize freq of intense convective events (related to Extreme Wx).